

# SOLUTIONS



1

## Definitions

A **solution** is a system in which one or more substances are *homogeneously* mixed or dissolved in another substance

- **homogeneous mixture**
  - uniform appearance
  - similar properties throughout mixture
- The **solvent** is the dissolving agent
  - *i.e.*, the most abundant component of the solution
- The **solute** is the component that is dissolved
  - *i.e.*, the least abundant component of the solution

2

## Concentration of solutions

**concentration** -- the amount of solute dissolved in a given quantity of solvent or solution

*There are many different types of concentration units:*

**molarity** ← *we will focus on this*

mass %

volume %

mass/volume %

parts per million (ppm)

parts per billion (ppb)

mole fraction

molality (not to be confused with *molarity*)

3

## Concentration units based on the number of moles of solute

**molarity** -- number of moles of *solute* per liter of *solution*

$$\text{Molarity} = \frac{\text{moles of solute}}{\text{liters solution}}$$

molarity has units of *moles per liter*  $\left(\frac{\text{mol}}{\text{L}}\right)$

which can be abbreviated as *M*



4

### Example: Preparation of a 1 molar solution of NaCl

1 mole NaCl = 58.44 g NaCl

1000.0-ml flask

water

$$\text{Molarity} = \frac{1.000 \text{ mol}}{1.000 \text{ L}} = 1.000 \text{ M}$$

fill line

Add 1 mole of NaCl to an empty 1-liter volumetric flask

Add water to completely dissolve NaCl

Add water until the fill mark is reached and mix thoroughly

5

### Example: Preparation of a 1 molar solution of NaCl

**Note:** Dissolving 1 mole of solute to make 1 liter of solution is **not** the only way to prepare a solution with a concentration of 1 M ( i.e., 1 mol / L )

0.250 mole NaCl = 14.61 g NaCl

250.0-ml flask

water

$$\text{Molarity} = \frac{0.2500 \text{ mol}}{0.2500 \text{ L}} = 1.000 \text{ M}$$

fill line

Add 0.250 mole of NaCl to an empty 250-ml volumetric flask

Add water to completely dissolve NaCl

Add water until the fill mark is reached and mix thoroughly

6

### Example: Preparation of a 0.5 molar solution of NaCl

0.5 mole NaCl = 29.22 g NaCl

1000.0-ml flask

fill line

$$\text{Molarity} = \frac{0.5000 \text{ mol}}{1.000 \text{ L}} = 0.5000 \text{ M}$$

0.250 mole NaCl = 14.61 g NaCl

500.0-ml flask

fill line

$$\text{Molarity} = \frac{0.2500 \text{ mol}}{0.5000 \text{ L}} = 0.5000 \text{ M}$$

7

**What is the molarity of a solution made by dissolving 2.00 g of potassium chlorate in enough water to make 150. ml of solution?**

**Step 1: Start with the definition of molarity:**

$$\text{Molarity} = \frac{\text{moles of solute}}{\text{liters solution}}$$

**Step 2: Determine the number of moles of solute**

$$\text{Molar mass of KClO}_3 = 39.10 + 35.45 + 3(16.00) = 122.6 \text{ g / mol}$$

$$2.00 \text{ g KClO}_3 \left( \frac{1 \text{ mole KClO}_3}{122.6 \text{ g KClO}_3} \right) = 0.0163 \text{ moles KClO}_3$$

8

What is the molarity of a solution made by dissolving 2.00 g of potassium chlorate in enough water to make 150. ml of solution?

$$\text{Molarity} = \frac{\text{moles of solute}}{\text{liters solution}}$$

Step 3: Determine the number of *liters* of solution

$$150. \cancel{\text{ml}} \left( \frac{1 \text{ liter}}{1000. \cancel{\text{ml}}} \right) = 0.150 \text{ L}$$

9

What is the molarity of a solution made by dissolving 2.00 g of potassium chlorate in enough water to make 150. ml of solution?

$$\text{Molarity} = \frac{\text{moles of solute}}{\text{liters solution}}$$

Step 4: Plug values into molarity equation

$$\text{Molarity} = \frac{0.0163 \text{ moles KClO}_3}{0.150 \text{ L}}$$

$$\text{Molarity} = 0.109 \text{ moles KClO}_3 / \text{L} = 0.109 \text{ M KClO}_3$$

10

How many grams of potassium hydroxide are required to prepare 500. ml of 0.450 M KOH solution?

Step 1: Start with the definition of molarity:

$$\text{Molarity} = \frac{\text{moles of solute}}{\text{liters solution}}$$

Step 2: Determine the number of *liters* of solution

$$500. \cancel{\text{ml}} \left( \frac{1 \text{ liter}}{1000. \cancel{\text{ml}}} \right) = 0.500 \text{ L}$$

11

How many grams of potassium hydroxide are required to prepare 500. ml of 0.450 M KOH solution?

$$\text{Molarity} = \frac{\text{moles of solute}}{\text{liters solution}}$$

Step 3: Plug known values into molarity equation and solve for unknown (moles of solute)

$$0.450 \text{ M KOH} = \frac{x \text{ moles of KOH}}{0.500 \text{ L}}$$

$$(\cancel{0.500 \text{ L}}) 0.450 \frac{\text{moles KOH}}{\cancel{\text{L}}} = \frac{x \text{ moles of KOH}}{\cancel{0.500 \text{ L}}} (\cancel{0.500 \text{ L}})$$

$$0.225 \text{ moles KOH} = x$$

12

How many grams of potassium hydroxide are required to prepare 500. ml of 0.450 M KOH solution?

$$\text{Molarity} = \frac{\text{moles of solute}}{\text{liters solution}}$$

Step 4: Convert moles KOH to grams KOH

$$\text{Molar mass of KOH} = 39.10 + 16.00 + 1.008 = 56.11 \text{ g/mol}$$

$$0.225 \text{ moles KOH} \left( \frac{56.11 \text{ g KOH}}{1 \text{ mole KOH}} \right) = 12.6 \text{ g KOH}$$

13

Calculate the number of moles of nitric acid in 325 ml of 16 M HNO<sub>3</sub>

Step 1: Start with the definition of molarity:

$$\text{Molarity} = \frac{\text{moles of solute}}{\text{liters solution}}$$

Step 2: Plug known values into molarity equation and solve for unknown (moles of solute)

$$16 \text{ M HNO}_3 = \frac{x \text{ moles of HNO}_3}{0.325 \text{ L}}$$

$$(0.325 \text{ L}) 16 \frac{\text{moles HNO}_3}{\cancel{\text{L}}} = \frac{x \text{ moles of HNO}_3}{\cancel{0.325 \text{ L}}} (0.325 \text{ L})$$

$$5.2 \text{ moles HNO}_3 = x$$

14

## Concentrations of ions in aqueous solutions


When an *ionic compound* dissolves, the concentrations of the individual ions depend on the chemical formula of the compound

**Example:** What are the concentrations of Na<sup>+</sup> and Cl<sup>-</sup> ions in a 0.125 M aqueous solution of NaCl?

$$0.125 \text{ M NaCl} = \frac{0.125 \text{ mol NaCl}}{\text{L solution}} \rightarrow \frac{0.125 \text{ mol Na}^+}{\text{L solution}} = 0.125 \text{ M Na}^+$$

$$\frac{0.125 \text{ mol Cl}^-}{\text{L solution}} = 0.125 \text{ M Cl}^-$$

Each NaCl formula unit produces 1 Na<sup>+</sup> ion and 1 Cl<sup>-</sup> ion in solution



15

## Concentrations of ions in aqueous solutions


When an *ionic compound* dissolves, the concentrations of the individual ions depend on the chemical formula of the compound

**Example:** What are the concentrations of Mg<sup>2+</sup> and Cl<sup>-</sup> ions in a 0.125 M aqueous solution of MgCl<sub>2</sub>?

$$0.125 \text{ M MgCl}_2 = \frac{0.125 \text{ mol MgCl}_2}{\text{L solution}} \rightarrow \frac{0.125 \text{ mol Mg}^{2+}}{\text{L solution}} = 0.125 \text{ M Mg}^{2+}$$

$$\frac{0.250 \text{ mol Cl}^-}{\text{L solution}} = 0.250 \text{ M Cl}^-$$

Each MgCl<sub>2</sub> formula unit produces 1 Mg<sup>2+</sup> ion and 2 Cl<sup>-</sup> ions in solution



16

A solution is made by dissolving 9.82 g of copper (II) chloride (CuCl<sub>2</sub>) in enough water to make 600. mL of solution. What is the molarity of Cl<sup>-</sup> ions in solution?

**Step 1: Determine the number of moles of solute**

Molar mass of CuCl<sub>2</sub> = 63.55 + 2(35.45) = **134.45 g / mol**

$$9.82 \text{ g } \cancel{\text{CuCl}_2} \left( \frac{1 \text{ mole CuCl}_2}{134.45 \text{ g } \cancel{\text{CuCl}_2}} \right) = \mathbf{0.0730 \text{ moles CuCl}_2}$$

**Step 2: Determine molarity of solute**


$$\text{Molarity} = \frac{0.0730 \text{ moles CuCl}_2}{0.600 \text{ L}} = \mathbf{0.122 \text{ M CuCl}_2}$$

17

A solution is made by dissolving 9.82 g of copper (II) chloride (CuCl<sub>2</sub>) in enough water to make 600. mL of solution. What is the molarity of Cl<sup>-</sup> ions in solution?

**Step 3: Determine the ion to solute ratio**

CuCl<sub>2</sub> dissociates to give one Cu<sup>2+</sup> ion and two Cl<sup>-</sup> ions



**Step 4: Determine molarity of the ion**

$$0.122 \text{ M CuCl}_2 \left( \frac{2 \text{ moles Cl}^- \text{ ions}}{1 \text{ mole CuCl}_2} \right) = \mathbf{0.244 \text{ M Cl}^- \text{ ions}}$$

18

## Dilutions

**Dilution:** Reducing the concentration of a solution by adding more solvent to the solution

- More solvent is added:  
-- *volume of the solution* increases
- No additional solute is added  
-- *number of moles of solute* stays the same

**Net result:** The *molarity* of the solution decreases

$$\downarrow \text{ Molarity} = \frac{\text{moles of solute (unchanged)}}{\text{liters solution } \uparrow}$$

19

● NO<sub>3</sub><sup>-</sup>

● Na<sup>+</sup>

**Moles = 1.0 mol**  
**Volume = 1.0 L**

$$\text{Molarity} = \frac{1.0 \text{ mol}}{1.0 \text{ L}} = \mathbf{1.0 \text{ M}}$$

**NaNO<sub>3</sub> solution**

20




Moles = 1.0 mol

Volume = 2.0 L

$$\text{Molarity} = \frac{1.0 \text{ mol}}{2.0 \text{ L}}$$

$$= 0.5 \text{ M}$$



- Solution volume is doubled
- Moles of solute remain the same
- Solution concentration is halved

21

Calculate the molarity of a solution prepared by diluting 125 ml of 0.400 M HCl to a final solution volume of 1.00 L.

For any dilution problem, remember that the number of moles of solute *remains the same*:

$$\text{moles of solute (before)} = \text{moles of solute (after)}$$

Based on the definition of molarity, this can be expressed as:

$$M_1 V_1 = M_2 V_2$$

Where  $M_1$  is the molarity of the original solution

$V_1$  is the volume of the original solution

$M_2$  is the molarity of the diluted solution

$V_2$  is the volume of the diluted solution

22

Calculate the molarity of a solution prepared by diluting 125 ml of 0.400 M HCl to a final solution volume of 1.00 L.

$$M_1 V_1 = M_2 V_2$$

$$M_1 = 0.400 \text{ M}$$

$$M_2 = ?$$

$$V_1 = 125 \text{ ml} = 0.125 \text{ L}$$

$$V_2 = 1.00 \text{ L}$$

$$\frac{(0.400 \text{ M})(0.125 \cancel{\text{ L}})}{1.00 \cancel{\text{ L}}} = \frac{(M_2)(1.00 \cancel{\text{ L}})}{1.00 \cancel{\text{ L}}}$$

$$0.0500 \text{ M} = M_2$$


23

## Solubility

**Solubility** refers to the ability of a compound to dissolve in a solvent  
 -- different compounds will dissolve to different extents in a given solvent


When water is the solvent:

If a solute dissolves readily in water, it is said to be **soluble** in water



Examples:  
 $\text{NaCl}$   
 $\text{AgNO}_3$   
 $(\text{NH}_4)_2\text{CO}_3$

If a solute will not dissolve in water, it is said to be **insoluble** in water



Examples:  
 $\text{Fe}(\text{OH})_3$   
 $\text{PbCl}_2$   
 $\text{CaCO}_3$

24

## Solubility rules

AN IONIC COMPOUND IS SOLUBLE IN WATER IF IT CONTAINS THE FOLLOWING IONS:	EXCEPTIONS
Ammonium ion ( $\text{NH}_4^+$ )	none
Alkali metal (Group IA) ions ( $\text{Li}^+$ , $\text{Na}^+$ , $\text{K}^+$ )	none
Nitrate ( $\text{NO}_3^-$ ) Acetate ( $\text{C}_2\text{H}_3\text{O}_2^-$ )	none
Halides ( $\text{Cl}^-$ , $\text{Br}^-$ , $\text{I}^-$ )	Compounds containing $\text{Ag}^+$ , $\text{Pb}^{2+}$ , $\text{Hg}_2^{2+}$
Sulfate ( $\text{SO}_4^{2-}$ )	Compounds containing $\text{Ag}^+$ , $\text{Pb}^{2+}$ , $\text{Ca}^{2+}$ , $\text{Sr}^{2+}$ , $\text{Ba}^{2+}$

25

## Solubility rules

AN IONIC COMPOUND IS NOT SOLUBLE IN WATER IF IT CONTAINS THE FOLLOWING IONS:	EXCEPTIONS
Carbonate ( $\text{CO}_3^{2-}$ ) Phosphate ( $\text{PO}_4^{3-}$ )	Compounds containing $\text{Li}^+$ , $\text{Na}^+$ , $\text{K}^+$ , $\text{NH}_4^+$ (soluble)
Hydroxide ( $\text{OH}^-$ )	Compounds containing $\text{Li}^+$ , $\text{Na}^+$ , $\text{K}^+$ , $\text{NH}_4^+$ (soluble) Compounds containing $\text{Ca}^{2+}$ , $\text{Ba}^{2+}$ , $\text{Sr}^{2+}$ (slightly soluble)
Sulfide ( $\text{S}^{2-}$ )	Compounds containing $\text{Li}^+$ , $\text{Na}^+$ , $\text{K}^+$ , $\text{NH}_4^+$ (soluble) Compounds containing $\text{Ca}^{2+}$ , $\text{Ba}^{2+}$ , $\text{Sr}^{2+}$ (soluble)

26

## Sample problems

Are the following compounds soluble or insoluble in water?

$\text{NaCl}$       **soluble**

$(\text{NH}_4)_3\text{PO}_4$       **soluble**

$\text{CaCO}_3$       **insoluble**

$\text{MgSO}_4$       **soluble**



$\text{BaSO}_4$       **insoluble**

27

## Double-displacement reactions

**double-displacement reaction** -- two ionic compounds exchange partners (*i.e.*, cations and anions) to form two different compounds

General form:




**Precipitation reactions** are a type of double-displacement reaction

28

## Reactions of aqueous solutions: *Precipitation reactions*

**Precipitation reactions** are a type of double-displacement reaction

In a **precipitation reaction**, an insoluble solid (called a precipitate) is formed when reactants in aqueous solution (*i.e.*, dissolved in water) are combined





↑  
insoluble precipitate  
indicated by (s) after its formula

29

## Precipitation reactions

Most **precipitation reactions** occur when the anions and cations of two aqueous ionic compounds switch partners


General form:



To predict whether a precipitation reaction will occur:

- look at the potential products of the reaction (*i.e.*, make the anions and cations switch partners)
- determine whether either product is an insoluble solid


Example: Will a precipitation reaction occur when aqueous zinc chloride and potassium hydroxide are mixed?




Use solubility rules to determine if either of these is an insoluble solid

30


## Will the following reactions take place?



↑  
insoluble precipitate



no reaction



↑  
insoluble precipitate

31

## For chemical reactions involving aqueous solutions, three types of equations can be written

### Molecular equation


Formulas written for all reactants & products do not show their ionic character -- *i.e.*, aqueous substances are shown as neutral compounds



### Complete ionic equation

All soluble strong electrolytes are shown as ions


-- aqueous substances are shown as separate cations and anions



### Net ionic equation

Includes only the substances that undergo change

-- ions that are present but do not react (spectator ions) are not shown




32




## Procedure for writing net ionic equations


### 1. Write a balanced molecular equation for the reaction



### 2. Rewrite equation to show aqueous substances as separate cations and anions (i.e., complete ionic equation)




### 3. Rewrite equation after identifying and canceling spectator ions



33

## Electrolytes

**electrolyte** -- a substance that forms ions when dissolved in water, resulting in a solution that conducts electricity



Electrolytes are capable of producing charge carriers (i.e., ions) in solution


Pure water

Non-electrolyte

34

## Electrolytes

**electrolyte** -- a substance that forms ions when dissolved in water, resulting in a solution that conducts electricity



Electrolytes are capable of producing charge carriers (i.e., ions) in solution



### Solution of NaCl in water

- presence of  $\text{Na}^{+}$  and  $\text{Cl}^{-}$  ions

NaCl is an **electrolyte**

35

## Formation of ions in solution



**Electrolytes** -- substances that form aqueous solutions containing ions

**Non-electrolytes** -- substances that do not form ions in solution

**dissociation** -- the separation of an ionic compound into its cations and anions as the compound dissolves


Example: Sodium chloride



36

## Electrolytes

**electrolyte** -- a substance that forms ions when dissolved in water, resulting in a solution that conducts electricity



37

## Strong and weak electrolytes

**Strong electrolytes** are solutes that exist in solution *completely or nearly completely* as ions

**Weak electrolytes** are solutes that dissociate only *partially* to form ions in solution

-- exist primarily as non-dissociated molecules in solution, with only a *small fraction* in the form of ions

- Nearly all *soluble ionic compounds* are strong electrolytes
- *Strong acids and bases* are strong electrolytes
- *Weak acids and bases* are weak electrolytes

We will talk about strong/weak acids and bases shortly

38